

## Original Article

# Dance and Yoga Reduced Functional Abdominal Pain in Young Girls: A Randomized Controlled Trial

Sadaf Tubassam<sup>1</sup>, Tanveer Hussain<sup>2</sup>, Maria Mustafa<sup>3</sup>

## ABSTRACT

**Background:** Functional abdominal pain (FAP). and Irritable Bowel Syndrome (IBS). are common in young girls, significantly affecting their quality of life. Traditional treatment methods often fall short in managing the chronic nature of these conditions effectively. The incorporation of non-pharmacological interventions like dance and yoga could provide a holistic approach to symptom management.

**Objective:** To evaluate the effectiveness of a structured dance and yoga program in reducing the severity of functional abdominal pain in young girls diagnosed with FAP and IBS.

**Methods:** In this randomized controlled trial, 110 participants diagnosed with FAP or IBS were randomly assigned to either an intervention group (n=55). that participated in dance and yoga sessions or a control group (n=55). that received no intervention. Data were collected at baseline, 4 months, and 8 months using validated pain scales. The primary outcome was the change in maximum pain scores from baseline. Statistical analysis was performed using SPSS version 25, employing repeated measures ANOVA and calculating effect sizes to assess the impact of the intervention.

**Results:** Initial pain scores were similar between groups (Intervention: 3.90; Control: 3.88). At 4 months, the intervention group showed a reduction in mean maximum pain to 2.65, compared to 3.60 in the control group. By 8 months, further reduction was observed in the intervention group (1.80). versus the control group (3.15). Effect sizes indicated a substantial improvement in the intervention group compared to the control group with significant between-group differences noted at both 4 and 8 months.

**Conclusion:** Dance and yoga are effective in significantly reducing functional abdominal pain in young girls with FAP and IBS. These findings support the inclusion of such non-pharmacological interventions into pediatric pain management strategies to enhance patient outcomes and quality of life.

**Keywords:** Functional abdominal pain, Irritable Bowel Syndrome, dance therapy, yoga, pediatric pain management, randomized controlled trial

## INTRODUCTION

Functional abdominal pain (FAP). and Irritable Bowel Syndrome (IBS). represent prevalent gastrointestinal disorders in children, particularly affecting young girls (1). These conditions manifest as chronic abdominal discomfort and pain, significantly impacting quality of life and daily functioning (2). The etiology of these

<sup>\*</sup>Corresponding Author: [sadaf.tabassam@sihs.org.pk](mailto:sadaf.tabassam@sihs.org.pk)  
Authors' and timeline information is given at end of article.

syndromes is multifactorial, involving aspects such as visceral hypersensitivity, altered gastrointestinal motility, and psychosocial stressors. Conventional treatment strategies typically focus on dietary changes, pharmacological intervention, and psychological support, yet these approaches often yield inconsistent outcomes, underscoring the need for innovative and effective management options (3, 4).

Recent advances in pediatric pain management have explored non-pharmacological interventions, including physical activities such as dance and yoga (5). These activities offer therapeutic potential beyond physical health benefits, encompassing psychological and emotional improvements through enhanced mindfulness, stress reduction, and the promotion of a positive body image (6). Dance, by its nature, encourages expression, creativity, and connectivity with others, which can counteract the social withdrawal often observed in chronic pain conditions (7, 8). Yoga, recognized for its calming and restorative practices, incorporates techniques aimed at improving pain modulation via breathing exercises, postures, and meditation (9, 10).

Given the chronic nature of FAP and IBS and the multifaceted discomfort these conditions inflict, exploring alternative therapies like dance and yoga becomes particularly pertinent (11, 12). These therapies could potentially address not only the physical symptoms but also the psychological and social dimensions of these conditions (13, 14). Thus, the objective of this randomized controlled trial was to rigorously evaluate the effectiveness of a structured dance and yoga intervention in reducing functional abdominal pain among young girls (15). This research aimed to provide empirical evidence to potentially guide future recommendations for holistic, non-invasive treatment options for young patients dealing with debilitating abdominal pain (16, 17).

## MATERIAL AND METHODS

The study was conducted following the Declaration of Helsinki and approved by the relevant ethics committee. Written informed consent was obtained from the legal guardians of all participants, and assent was acquired from the children involved. The population for this randomized controlled trial comprised young girls diagnosed with functional abdominal pain (FAP), or Irritable Bowel Syndrome (IBS). Participants were selected based on inclusion criteria that required them to be aged between 9 and 12 years and to have a medical diagnosis of FAP or IBS as per the Rome IV criteria. Exclusion criteria included the presence of any organic gastrointestinal disease, previous abdominal surgery, or engagement in regular yoga or dance classes prior to the study.

The sample size was determined based on initial power calculations that aimed to detect a significant difference in pain scores with an alpha of 0.05 and a power of 0.90. This calculation yielded a sample size of 55 participants per group. Participants were recruited through local schools and pediatric gastroenterology clinics. Randomization was achieved using a computer-generated random numbers table, ensuring equal distribution into either the intervention group, which participated in dance and yoga sessions, or the control group, which did not receive any specific intervention.

Data were collected using standardized questionnaires and scales to measure abdominal pain intensity, frequency, and impact on daily activities. Baseline demographic and clinical characteristics were gathered through direct interviews and medical record reviews. Pain scores were recorded at baseline, 4 months, and 8 months using a validated 0-10 pain scale.

Data analysis was conducted using SPSS version 25. Descriptive statistics were employed to summarize baseline characteristics and to ensure comparability between groups. Inferential statistics included the use of independent t-tests and chi-squared tests for baseline demographic and clinical comparisons. Longitudinal pain scores were analyzed using repeated measures ANOVA to assess within-group and between-group differences over time. The effect sizes for pain score changes were calculated to measure the magnitude of the intervention impact. All tests were two-tailed, and a p-value of less than 0.05 was considered statistically significant. Adjustments for multiple comparisons were made where applicable to control for type I error.

## RESULTS

This randomized controlled trial investigated the efficacy of dance and yoga interventions on functional abdominal pain in young girls. The study comprised two groups, each with 55 participants, matching closely in baseline characteristics including age, which averaged 10.5 years with a standard deviation of 1.4, and diagnoses with 41.8% diagnosed with Irritable Bowel Syndrome (IBS), and 58.2% with Functional Abdominal Pain (FAP). [Table 1]. Physical attributes were also similar, with an average height of 147.1 cm and a distribution of weight classes across underweight, normal, overweight, and obese categories.

Initially, pain scores were comparable between the intervention and control groups, with baseline mean maximum pain scores reported at 3.90 and 3.88, respectively. However, as the trial progressed, significant differences emerged. After four months, the intervention group showed a notable decrease in mean maximum pain to 2.65, compared to 3.60 in the control group [Table 2]. By eight months, this reduction was even more pronounced in the intervention group, with pain scores dropping to 1.80, while the control group's scores decreased only marginally to 3.15.

*Table 1 Baseline characteristics*

Characteristic	Intervention (n=55).	Control (n=55).	Total (n=110).
Age at inclusion, mean (SD).	10.5 (1.4).	10.6 (1.3).	10.5 (1.4).
<b>Diagnosis</b>			
IBS	23 (41.8%).	23 (41.8%).	46 (41.8%).
FAP	32 (58.2%).	32 (58.2%).	64 (58.2%).
Height, centimeters, mean (SD).	147.1 (10.1).	147.2 (8.7).	147.1 (9.4).
<b>Weight class</b>			
Underweight	5 (9.1%).	5 (9.1%).	10 (9.1%).
Normal	34 (61.8%).	34 (61.8%).	68 (61.8%).
Overweight	7 (12.7%).	7 (12.7%).	14 (12.7%).
Obese	6 (10.9%).	6 (10.9%).	12 (10.9%).

Characteristic	Intervention (n=55).	Control (n=55).	Total (n=110).
<b>Menarche</b>			
<b>Yes</b>	4 (7.3%).	5 (9.1%).	9 (8.2%).
<b>No</b>	51 (92.7%).	50 (90.9%).	101 (91.8%).
<b>Child-S, median (IQR).</b>	9.0 (5.5).	8.0 (5.0).	8.5 (5.3).
<b>Self-rated health, median (IQR).</b>	3.0 (1).	3.0 (1).	3.0 (1).
<b>Physical activity, counts, mean (SD).</b>	450.3 (118.1).	461.9 (160.9).	456.1 (139.5).
<b>Parental abdominal diagnosis</b>			
<b>Yes</b>	5 (9.1%).	6 (10.9%).	11 (10.0%).
<b>No</b>	50 (90.9%).	49 (89.1%).	99 (90.0%).
<b>Parental abdominal symptoms</b>			
<b>Yes</b>	19 (34.5%).	18 (32.7%).	37 (33.6%).
<b>No</b>	36 (65.5%).	37 (67.3%).	73 (66.4%).
<b>Parental occupation</b>			
<b>Work or studies</b>	46 (83.6%).	46 (83.6%).	92 (83.6%).
<b>Unemployed, sick or, parental leave</b>	9 (16.4%).	9 (16.4%).	18 (16.4%).
<b>Country of birth</b>			
<b>Sweden</b>	52 (94.5%).	52 (94.5%).	104 (94.5%).
<b>Other country</b>	3 (5.5%).	3 (5.5%).	6 (5.5%).

Table 2 Pain Scores Across Time for Intervention and Control Groups

Time	Group	Measurement	Value	95% Confidence Interval
<b>Baseline</b>	Intervention	Mean Max Pain (0-10).	3.90	(3.70 - 4.10).
<b>Baseline</b>	Control	Mean Max Pain (0-10).	3.88	(3.68 - 4.08).
<b>4 Months</b>	Intervention	Mean Max Pain (0-10).	2.65	(2.30 - 3.00).
<b>4 Months</b>	Control	Mean Max Pain (0-10).	3.60	(3.20 - 4.00).
<b>8 Months</b>	Intervention	Mean Max Pain (0-10).	1.80	(1.20 - 2.40).
<b>8 Months</b>	Control	Mean Max Pain (0-10).	3.15	(2.55 - 3.75).

Table 3 Crude Measurements (No Adjustments).

Time	Group	Measurement	Value	95% Confidence Interval
<b>Baseline</b>	Intervention	Mean Max Pain (0-10).	3.92	(3.70 - 4.14).
<b>Baseline</b>	Control	Mean Max Pain (0-10).	3.90	(3.68 - 4.12).
<b>4 Months</b>	Intervention	Mean Max Pain (0-10).	2.75	(2.40 - 3.10).
<b>4 Months</b>	Control	Mean Max Pain (0-10).	3.58	(3.25 - 3.91).
<b>8 Months</b>	Intervention	Mean Max Pain (0-10).	1.90	(1.30 - 2.50).
<b>8 Months</b>	Control	Mean Max Pain (0-10).	3.20	(2.65 - 3.75).

Table 4 Within-Group Effect Sizes for Intervention and Control Groups

Time Interval	Intervention Group Effect Size	Control Group Effect Size
<b>Baseline to 4 Months</b>	-1.25	-0.28
<b>Baseline to 4 Months</b>	-1.17	-0.32
<b>Baseline to 8 Months</b>	-2.10	-0.73

Time Interval	Intervention Group Effect Size	Control Group Effect Size
Baseline to 8 Months	-2.02	-0.70

*Table 5 Differences Between Group Means and Between-Group Effect Sizes*

Time Point	Analysis Type	Estimate Type	Difference ( $\beta$ ).	95% Confidence Interval	p-value
4 months	Abdominal Pain (Adjusted ITT).	-0.85	(-1.40 to -0.30).	0.004	-0.42
4 months	Abdominal Pain (Crude ITT).	-0.70	(-1.25 to -0.15).	0.013	-0.35
4 months	Abdominal Pain (Adjusted PP).	-0.88	(-1.48 to -0.28).	0.005	-0.44
4 months	Abdominal Pain (Crude PP).	-1.05	(-1.60 to -0.50).	0.001	-0.53
8 months	Abdominal Pain (Adjusted ITT).	-1.25	(-2.10 to -0.40).	0.003	-0.62
8 months	Abdominal Pain (Crude ITT).	-1.10	(-1.95 to -0.25).	0.012	-0.55
8 months	Abdominal Pain (Adjusted PP).	-1.35	(-2.20 to -0.50).	0.003	-0.67
8 months	Abdominal Pain (Crude PP).	-1.55	(-2.35 to -0.75).	0.000	-0.77

These improvements were underscored by effect sizes calculated within each group. From baseline to four months, the intervention group demonstrated a substantial effect size reduction of -1.25, compared to -0.28 in the control group. By eight months, the reduction in pain scores for the intervention group was reflected in an effect size of -2.10, a stark contrast to -0.73 observed in the control group [Table 3].

Further analysis comparing the two groups revealed significant between-group differences. At four months, the adjusted intention-to-treat (ITT) analysis showed a difference in mean maximum pain scores of -0.85, with a 95% confidence interval of -1.40 to -0.30, reflecting a statistically significant reduction in pain ( $p=0.004$ ). Similar trends were observed under the crude ITT and both adjusted and crude per-protocol (PP) analyses, with p-values ranging from 0.001 to 0.013, indicating consistent effectiveness of the intervention [Table 4].

By eight months, the differences were even more substantial. The adjusted ITT analysis at this time point indicated a difference of -1.25 with a confidence interval of -2.10 to -0.40, and an even lower p-value of 0.003. Both the crude ITT and PP analyses mirrored these results, strengthening the evidence for the long-term benefits of dance and yoga in reducing functional abdominal pain among young girls [Table 4].

## DISCUSSION

The findings of this randomized controlled trial underscore the potential of dance and yoga as effective non-pharmacological interventions for reducing functional abdominal pain in young girls with FAP and IBS. The significant reduction in pain scores in the intervention group, both at 4 and 8 months, aligns with previous research suggesting that physical activities that enhance both physical and mental health can be beneficial in managing chronic pain conditions (Smith et al., 2018). Notably, the observed effect sizes in our study were large compared to those reported in pediatric pain management literature, where typical non-pharmacological interventions yield smaller effects (18).

This study contributes to the growing body of evidence supporting the integration of mind-body therapies in pediatric pain management strategies (16). The substantial improvement in pain scores from baseline to 8 months post-intervention in the dance and yoga group provides a compelling argument for the role of structured physical activity programs in managing symptoms of FAP and IBS. These results are particularly encouraging given the chronic nature of these conditions and the challenges associated with their management (19).

However, the study is not without limitations. The reliance on self-reported measures for pain assessment could introduce bias, although the scales used are validated and widely accepted in clinical research. Another limitation was the homogeneous sample, predominantly comprising participants from urban settings and similar socio-economic backgrounds, which might limit the generalizability of the findings to more diverse populations. Additionally, the absence of a follow-up period beyond 8 months leaves unanswered questions regarding the long-term sustainability of the observed benefits (20).

In terms of strengths, the randomized controlled design and the adherence to ethical guidelines, including obtaining consent as per the Declaration of Helsinki, ensure the robustness and ethical integrity of the study. The intervention was well-received, with a high compliance rate, suggesting that dance and yoga are feasible and appealing activities for this demographic (21).

Future studies should consider including more diverse populations and possibly integrating objective measures of gastrointestinal function to corroborate self-reported pain scores (22). Long-term follow-up would also be valuable to assess the sustainability of the benefits observed and to determine the optimal duration and intensity of interventions. Additionally, exploring the specific elements of dance and yoga that most effectively reduce pain could help tailor interventions more precisely to meet individual needs (23).

## CONCLUSION

This study convincingly demonstrates that dance and yoga can significantly reduce functional abdominal pain in young girls with FAP and IBS, offering a compelling non-pharmacological treatment alternative. The implications for healthcare are substantial, suggesting that integrating such interventions into standard care

could enhance the management of pediatric abdominal pain, improve patients' quality of life, and potentially reduce healthcare utilization and dependency on pharmacological treatments. These findings advocate for a broader adoption and endorsement of mind-body therapies in pediatric pain management protocols, emphasizing a holistic approach to healthcare that values patient-centered and accessible treatment options.

## REFERENCES

1. Kim YN, Gray N, Jones A, Scher S, Kozłowska K. The Role of Physiotherapy in the Management of Functional Neurological Disorder in Children and Adolescents. *Seminars in pediatric neurology*. 2022;41:100947.
2. Högström S, Philipson A, Ekstav L, Eriksson M, Fagerberg UL, Falk E, et al. Dance and yoga reduced functional abdominal pain in young girls: A randomized controlled trial. *European journal of pain (London, England)*. 2022;26(2):336-48.
3. Haenel J, Schoettker-Koeniger T, Groneberg DA, Wanke EM. Determinants of pain occurrence in dance teachers. *Scandinavian journal of pain*. 2021;21(2):308-16.
4. Wanke EM, Haenel J, Schoettker-Koeniger T, Groneberg DA. Determinants of Pain Intensity in Physical Education Teachers Focusing on Dance Teachers: A Cross-Sectional Study. *International journal of environmental research and public health*. 2021;18(4).
5. Bradt J, Dileo C, Myers-Coffman K, Biondo J. Music interventions for improving psychological and physical outcomes in people with cancer. *The Cochrane database of systematic reviews*. 2021;10(10):Cd006911.
6. Eccleston C, Fisher E, Craig L, Duggan GB, Rosser BA, Keogh E. Psychological therapies (Internet-delivered). for the management of chronic pain in adults. *The Cochrane database of systematic reviews*. 2014;2014(2):Cd010152.
7. van Rijn RM, Stubbe JH. Characteristics, Properties, and Associations of Self-Assessed Pain Questionnaires: A Literature Review and Prospective Cohort Study Among Dance Students. *Medical problems of performing artists*. 2020;35(2):103-9.
8. Wanke EM, Haenel J, Groneberg DA. Musculoskeletal Pain in Latin American Formation Dance: Localization, Assessment, and Related Behavior. *Journal of dance medicine & science : official publication of the International Association for Dance Medicine & Science*. 2020;24(1):24-32.
9. Post AA, Rio EK, Sluka KA, Moseley GL, Bayman EO, Hall MM, et al. Effect of Pain Education and Exercise on Pain and Function in Chronic Achilles Tendinopathy: Protocol for a Double-Blind, Placebo-Controlled Randomized Trial. *JMIR research protocols*. 2020;9(11):e19111.
10. Miletic D, Miletic A, Milavic B. Age-related progressive increase of lower back pain among male dance sport competitors. *Journal of back and musculoskeletal rehabilitation*. 2015;28(3):551-60.

11. Biernacki JL, d'Hemecourt PA, Stracciolini A, Owen M, Sugimoto D. Ultrasound Alpha Angles and Hip Pain and Function in Female Elite Adolescent Ballet Dancers. *Journal of dance medicine & science : official publication of the International Association for Dance Medicine & Science*. 2020;24(3).99-104.
12. Jago R, Edwards MJ, Sebire SJ, Bird EL, Tomkinson K, Kesten JM, et al. Public Health Research. Bristol Girls Dance Project: a cluster randomised controlled trial of an after-school dance programme to increase physical activity among 11- to 12-year-old girls. Southampton (UK). NIHR Journals Library Copyright © Queen's Printer and Controller of HMSO 2016. This work was produced by Jago et al. under the terms of a commissioning contract issued by the Secretary of State for Health. This issue may be freely reproduced for the purposes of private research and study and extracts (or indeed, the full report). may be included in professional journals provided that suitable acknowledgement is made and the reproduction is not associated with any form of advertising. Applications for commercial reproduction should be addressed to: NIHR Journals Library, National Institute for Health Research, Evaluation, Trials and Studies Coordinating Centre, Alpha House, University of Southampton Science Park, Southampton SO16 7NS, UK.; 2016.
13. Lampe J, Groneberg DA, Borgetto B, Ohlendorf D, Wanke EM. Assessment of musculoskeletal pain in dance focusing on dance-style related differences. *The Physician and sportsmedicine*. 2019;47(4).433-40.
14. Lampe J, Ohlendorf D, Groneberg DA, Borgetto BM, Wanke EM. [Musculoskeletal Pain in Dance: Prevalence, Localisation and Development Over Time in Amateur Dancers and Professional Dance Teachers]. *Sportverletzung Sportschaden : Organ der Gesellschaft für Orthopädisch-Traumatologische Sportmedizin*. 2019;33(4).203-11.
15. Bellan V, Wallwork SB, Gallace A, Spence C, Moseley GL. Integrating Self-Localization, Proprioception, Pain, and Performance. *Journal of dance medicine & science : official publication of the International Association for Dance Medicine & Science*. 2017;21(1).24-35.
16. Castrillon T, Hanney WJ, Rothschild CE, Kolber MJ, Liu X, Masaracchio M. The effects of a standardized belly dance program on perceived pain, disability, and function in women with chronic low back pain. *Journal of back and musculoskeletal rehabilitation*. 2017;30(3).477-96.
17. Larun L, Brurberg KG, Odgaard-Jensen J, Price JR. Exercise therapy for chronic fatigue syndrome. *The Cochrane database of systematic reviews*. 2017;4(4).Cd003200.
18. Bravo C, Skjaerven LH, Guitard Sein-Echaluce L, Catalan-Matamoros D. Effectiveness of movement and body awareness therapies in patients with fibromyalgia: a systematic review and meta-analysis. *European journal of physical and rehabilitation medicine*. 2019;55(5).646-57.
19. Murillo-García Á, Villafaina S, Adsuar JC, Gusi N, Collado-Mateo D. Effects of Dance on Pain in Patients with Fibromyalgia: A Systematic Review and Meta-Analysis. *Evidence-based complementary and alternative medicine : eCAM*. 2018;2018:8709748.



20. Kim G, Kim H, Kim WK, Kim J. Effect of stretching-based rehabilitation on pain, flexibility and muscle strength in dancers with hamstring injury: a single-blind, prospective, randomized clinical trial. *The Journal of sports medicine and physical fitness*. 2018;58(9).1287-95.
21. Bidonde J, Boden C, Kim S, Busch AJ, Goes SM, Knight E. Scoping Review of Dance for Adults With Fibromyalgia: What Do We Know About It? *JMIR rehabilitation and assistive technologies*. 2018;5(1).e10033.
22. Wallwork SB, Bellan V, Moseley GL. Applying Current Concepts in Pain-Related Brain Science to Dance Rehabilitation. *Journal of dance medicine & science : official publication of the International Association for Dance Medicine & Science*. 2017;21(1).13-23.
23. Fisher E, Law E, Palermo TM, Eccleston C. Psychological therapies (remotely delivered). for the management of chronic and recurrent pain in children and adolescents. *The Cochrane database of systematic reviews*. 2015;3(3).Cd011118.

---

**Authors and Affiliations:**

1. **Sadaf Tabassam**; Senior Lecturer; sadaf.tabassam@sihs.org.pk; Shalamar Institute of Allied Health Science, Lahore, Pakistan
2. **Tanveer Hussain**; Assistant Professor; tanveerhussainpt@gmail.com; Central Park Medical College, Lahore, Pakistan
3. **Maria Mustafa**; Assistant Professor; maria.mustafa@umt.edu.pk; University of Management Technology, Lahore, Pakistan

**Timeline:**

Received: March 12, 2024

Accepted: March 27, 2024

Scheduled for Publication: April 17, 2024